

COULD TH-234 PARTITIONING IN SEAWATER BE KRILL-DRIVEN?

Alessia-M. Rodriguez Y Baena¹, Scott-W. Fowler² * and Michel Warnau¹

¹ International Atomic Energy Agency, Marine Environment Laboratories, 4 Quai Antoine 1er, MC-98000 Monaco

² Marine Sciences Research Center, SUNY, Stony Brook, New York 11794-5000. - s.fowler@free.fr

Abstract

²³⁴Th is considered a useful tracer of oceanic biogeochemical processes occurring over timescales of days to weeks. Traditionally, ²³⁴Th-based particle export mathematical models have not taken into consideration the biological compartment, as typical zooplankton biomass values are not likely to result in significant biologically-mediated interactions with such models. However, recent experimental results for the Mediterranean krill *Meganyctiphanes norvegica* suggest that, if rapid ²³⁴Th uptake by krill is not considered in the modeling, the occurrence of a synchronized molting event within a high-biomass krill school may result in an overestimation of ²³⁴Th based particle flux by as much as one order of magnitude.

Keywords : *Crustacea, Models, Particle Flux, Radionuclides.*

Introduction

²³⁴Th is a naturally-occurring radionuclide constantly produced in seawater by its soluble parent ²³⁸U. Due to its high particle reactivity and relatively short half-life ($t_{1/2} = 24.1$ days), ²³⁴Th is commonly used as a proxy to estimate POC export from the upper oceanic water column [1]. This export is usually assessed by quantifying total ²³⁴Th deficits with respect to its conservative parent nuclide ²³⁸U in seawater (*ibid.*). Recent findings for three species of Antarctic crustaceans have suggested that under certain conditions krill-associated ²³⁴Th would generate a several-fold positive export bias in depth-integrated ²³⁴Th profiles [2]. Thus, we have tested this hypothesis by conducting analogous experiments using specimens of *Meganyctiphanes norvegica* krill collected in the NW Mediterranean.

Material and Methods

Live *Meganyctiphanes norvegica* from the NW Mediterranean were transported to the IAEA-MEL premises (Monaco), where they were acclimated for two weeks to laboratory conditions simulating their original temperature and light conditions (14°C; 37 p.s.u; darkness).

Specimens were individually placed in a 500 ml container containing 400 ml of natural filtered seawater spiked with ²³⁴Th. The seawater was changed and the radionuclide spike was renewed daily. During seawater renewal operations, animals were fed shortly on a mixture of uncontaminated phytoplankton and freshly hatched brine shrimp larvae.

At different times, each individual was γ -counted (high-resolution γ -spectrometry system) alive to determine the radionuclide uptake kinetics. At the end of the 7.5-d exposure period, the individuals were placed for 10 d in new containers with clean filtered seawater that was renewed daily. Individuals were then γ -counted daily to determine radionuclide loss kinetics.




Fig. 1. *Euphausia superba* - Krill (Edgar, 1997)

Results and Discussion

Krill rapidly took up ²³⁴Th, reaching a steady-state concentration factor of about 300 in less than one week, and strongly retained the accumulated radionuclide until molting occurred. At 14 °C krill molt on average every 8 days [3], and during the experiment 9 molting events were observed which indicated that ca. 50% of whole-body ²³⁴Th activity was associated with

the exoskeleton. Immediately following molting a corresponding amount of ²³⁴Th was rapidly adsorbed on the newly formed cuticle.

These results, coupled with published values of krill densities as high as 100 g l⁻¹ [4] (Fig. 1) which commonly occur in the Southern Ocean suggest that ²³⁴Th bioaccumulation could result in up to >90 % of total water-column ²³⁴Th being associated with krill (Fig. 2). Hence, the occurrence of a synchronized molting event [3] within a high-biomass krill school could cause the ²³⁴Th distribution in the water column to be almost entirely a result of radionuclide uptake by these organisms and consequently bias the results of traditional ²³⁴Th based models by more than one order of magnitude.




Fig. 2. CF-based computation representing the fraction (%) of total ²³⁴Th in the water column associated with different biomasses of crustacean zooplankton (g l⁻¹ wet wt), based on mean CF derived from experiments on Antarctic isopods and amphipods (fine dashed line [2]) or on Mediterranean krill (heavy solid line; present study).

References

- 1 - Waples J.T., Benitez-Nelson C., Savoye N., Rutgers van der Loeff M., Baskaran M. and O. Gustafsson, 2006. An introduction to the application and future use of ²³⁴Th in aquatic systems. *Mar. Chem.* 100: 166-189.
- 2 - Rodriguez y Baena A.M., Metian M., Teyssié J.L., De Broyer C. and M. Warnau, 2006. Experimental evidence for ²³⁴Th bioaccumulation in three Antarctic crustaceans: potential implications in particle flux studies. *Mar. Chem.* 100 (354-365)
- 3 - Hamner W.M., Hamner P.P., Strand S.W. and R.W.Gilmer, 1983. Behavior of Antarctic krill, *Euphausia superba*; chemoreception, feeding, schooling, and molting. *Science* 220: 433-435.
- 4 - Hewitt R.P., Watkins J., Naganobu M., Sushin V., Brierley A.S., Demer D., Kasatkina S., Takao Y., Goss C. and A. Malysenko, 2004. Biomass of Antarctic krill in the Scotia Sea in January/February 2000 and its use in revising an estimate of precautionary yield. *Deep Sea Res. Part II* 51: 1215-1236.